

Package ‘moonlit’

March 6, 2026

Type Package

Title Predicting Moonlight Intensity for a Given Time and Location

Version 0.1.1

Description Tools for predicting moonlight intensity on the ground based on the position of the moon, atmospheric conditions, and other factors. Provides functions to calculate moonlight intensity and related statistics for ecological and behavioral research, offering more accurate estimates than simple moon phase calculations. The underlying model is described in Smielak (2023) <[doi:10.1007/s00265-022-03287-2](https://doi.org/10.1007/s00265-022-03287-2)>.

License GPL-3

Encoding UTF-8

RoxygenNote 7.3.2

Depends R (>= 3.5.0)

Imports suncalc, lubridate

Suggests testthat (>= 3.0.0)

Config/testthat/edition 3

URL <https://github.com/msmielak/moonlit>

BugReports <https://github.com/msmielak/moonlit/issues>

NeedsCompilation no

Author Michal Smielak [aut, cre] (ORCID:
<<https://orcid.org/0000-0002-3585-9945>>)

Maintainer Michal Smielak <michalsmielak@gmail.com>

Repository CRAN

Date/Publication 2026-03-06 13:50:02 UTC

Contents

calculateMoonlightIntensity	2
calculateMoonlightStatistics	3
elevExtCoeff	4

calculateMoonlightIntensity
Calculate moonlight intensity

Description

This function predicts moonlight intensity on the ground for any given place and time, based on the location, position of the moon and number of correction factors

Usage

```
calculateMoonlightIntensity(lat, lon, date, e)
```

Arguments

lat	Latitude, numerical decimal
lon	Longitude, numerical decimal
date	Date time as POSIXct with the local time zone. If needed use as.POSIXct(date, tz=timezone)
e	Extinction coefficient - a single numerical value depending on the altitude. Average extinction coefficients (magnitude per air mass) are as follows: At sea level: 0.28; at 500m asl: 0.24; at 1000m asl: 0.21; at 2000m asl: 0.16

Value

A data frame with the following columns:

night	Logical, TRUE when sun is below the horizon
sunAltDegrees	Solar altitude in degrees
moonlightModel	Predicted moonlight illumination, relative to an "average" full moon
twilightModel	Predicted twilight illumination in lux
illumination	Combined moon and twilight intensity, in lux
moonPhase	Lunar phase as a numerical value (% of moon face illuminated)

Examples

```
lat <- 52.2297
lon <- 21.0122
date <- as.POSIXct("2023-06-15 22:00:00", tz = "Europe/Warsaw")
result <- calculateMoonlightIntensity(lat, lon, date, e = 0.21)
```

```
calculateMoonlightStatistics
```

Calculate nightly moonlight statistics

Description

Calculate nightly moonlight statistics

Usage

```
calculateMoonlightStatistics(lat, lon, date, e, t, timezone)
```

Arguments

lat	latitude, numerical decimal
lon	longitude, numerical decimal
date	date time as POSIXct with the local time zone. If needed use as.POSIXct(date, tz=timezone)
e	extinction coefficient - a single numerical value depending on the altitude. Average extinction coefficients (magnitude per air mass) are as follows: (At sea level: 0.28; at 500m asl: 0.24; at 1000m asl: 0.21; at 2000m asl: 0.16)
t	sampling interval. It is used in seq() function so the same values are accepted: A character string, containing one of "sec", "min", "hour". This can optionally be preceded by a (positive or negative) integer and a space, or followed by "s". Example: "15 mins", "3 hour" etc.
timezone	time zone of the data usually in the format "Continent/City", i.e. for Poland: "Europe/Warsaw"

Value

A data frame with one row per input record and the following columns:

sunset	Time of sunset for the night the record belongs to
sunrise	Time of sunrise for the night the record belongs to
date	Original date-time from the input
meanMoonlightIntensity	Mean nightly moonlight illumination
maxMoonlightIntensity	Maximum nightly moonlight illumination
minMoonlightIntensity	Minimum nightly moonlight illumination
meanMoonPhase	Mean nightly moon phase (fraction of disc illuminated)
maxMoonPhase	Maximum nightly moon phase
minMoonPhase	Minimum nightly moon phase

Examples

```
lat <- 52.2297
lon <- 21.0122
date <- as.POSIXct("2023-06-15 22:00:00", tz = "Europe/Warsaw")
result <- calculateMoonlightStatistics(lat, lon, date, e = 0.21,
                                     t = "30 mins",
                                     timezone = "Europe/Warsaw")
```

`elevExtCoeff`*Calculate extinction coefficient based on elevation of the observer*

Description

Calculate extinction coefficient based on elevation of the observer

Usage

```
elevExtCoeff(elev)
```

Arguments

`elev` elevation in meters asl

Value

A single numeric value: the estimated extinction coefficient for the given elevation.

Examples

```
# Extinction coefficient at sea level
suppressWarnings(elevExtCoeff(0))

# Extinction coefficient at 1000 m asl (e.g. approximate for many study sites)
suppressWarnings(elevExtCoeff(1000))
```

Index

calculateMoonlightIntensity, [2](#)
calculateMoonlightStatistics, [3](#)
elevExtCoeff, [4](#)